

Tularemia

Tularemia is a Class A disease. It must be reported to the state within 24 hours by calling the phone number listed on the website.

Tularemia is a zoonotic disease, transmitted from animals to humans. Originally described in the United States in 1911, the disease has been reported in all states except Hawaii. However, disease surveillance indicates a low level of natural transmission.


Tularemia is a bacterial disease caused by the gram-negative coccobacillus, *Francisella tularensis*. In the U.S., most people acquire the infection by arthropod bites - primarily ticks, or by contact with infected mammals - particularly rabbits. Deerflies can also serve as vectors and transmission from other animals, such as muskrats, is possible. Laboratory exposure, transmission through contaminated drinking water or food, and transmission via inhalation of contaminated dust or aerosols have also occurred.

The disease usually presents as an acute febrile illness. Several clinical presentations, depending on route of entry, can occur, including mucus membrane or skin ulceration, pharyngitis, ocular lesions, regional lymph-adenopathy and pneumonia.

Presumptive diagnosis can be made by serology or by the detection of antigen in tissues. Culture of the organism or demonstration of a four-fold increase in antibody titer is used for confirmation.

Reporting of tularemia was suspended in 1994, but was reintroduced in 2000 due to concerns with use of the agent as a biological weapon. In fact, outbreaks of pneumonic tularemia, especially in low-incidence areas, should initiate suspicion of intentional use. Only ten cases have been reported in Louisiana since 1987. (Figure 1)

Figure 1: Tularemia cases Louisiana, 1987-2011



Cases of tularemia that occur in summer are usually associated with arthropod bites, while winter exposures are commonly associated with hunter contact of rabbit carcasses. In recent years, a seasonal increase in late spring and summer has occurred when arthropod bites are most common. Nearly 70% of cases report an onset from May through August.